

U.S. DEPARTMENT OF THE INTERIOR  
U.S. GEOLOGICAL SURVEY

COMPILATION OF 29 SONIC AND DENSITY LOGS FROM 23 OIL TEST WELLS IN  
WESTERN WASHINGTON STATE

By

Thomas M. Brocher<sup>1</sup> and April L. Ruebel<sup>1</sup>

Open-File Report 98-249

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards or with the North American Stratigraphic Code. Any use of trade, product or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

<sup>1</sup>345 Middlefield Road, M/S 977, Menlo Park, CA 94025

1998

## **ABSTRACT**

Three-dimensional velocity models for Puget Sound provide a means for better understanding the lateral variations in strong ground motions recorded during local earthquakes in Puget Lowland. We have compiled 29 sonic and density logs from 23 oil test wells to help us determine the geometry and physical properties of the Cenozoic basins in western Washington. The maximum depths sampled by the test wells are between 0.47 and 4.04 km. These well logs sample Quaternary to Eocene sedimentary and volcanic rocks. This report presents the locations, elevations, depths, stratigraphic and other information about the oil test wells, and provides plots showing the density and sonic velocities as a function of depth for each well log. We also present two-way travel times calculated from the sonic velocities.

## CONTENTS

Abstract	1
Introduction	4
Well Log Analysis	5
Comments on the Well Log Data	6
Data Availability	7
Discussion	8
Acknowledgments	8
References	8
Abbreviations Used in Table 1	9

## TABLE

Table 1. Oil test well locations and drilling data	10
Table 2. Stratigraphy in the oil test wells	11
Table 3. Linear regression of the sonic velocity logs	13
Table 4. Linear regression of the density logs	14

## FIGURES

Figure 1. Map showing locations of oil test wells analyzed within this report	15
Figure 2. Sonic velocities for Kerry BN #34-11, L B Petroleum	16
Figure 3. Sonic velocities for Amoco-Weyerhaeuser #1-29, Amoco Production	17
Figure 4. Sonic velocities for Blessing Siler #1, Carr Edward J	18
Figure 5. Sonic velocities for Washington State #1, Phillips Petroleum Co.	19
Figure 6. Sonic velocities for Brandt #1, McCroskey A E Syndicate	20
Figure 7. Sonic velocities for Brandt #2, McCroskey A E Syndicate	21
Figure 8. Sonic velocities for KSD #1, Geothermal Resources Int'l.	22
Figure 9. Sonic velocities for Socal-Schroeder #1, Standard Oil of Calif.	23
Figure 10. Sonic velocities for Kingston #1, Mobil Oil	24
Figure 11. Sonic velocities for Pope & Talbot #18-1, Union Oil of Calif.	25
Figure 12. Sonic velocities for Socal Whidbey #1, Standard Oil of Calif.	26
Figure 13. Sonic velocities for Pope & Talbot #3-1, Standard Oil of Calif.	27
Figure 14. Sonic velocities for Dungeness #1, Standard Oil of Calif.	28
Figure 15. Sonic velocities for R D Merrill Co. #1, Texaco Inc.	29
Figure 16. Sonic velocities for Silvana Community #12-1, Standard Oil of Calif.	30
Figure 17. Sonic velocities for Merrill-Ring #1, Russell A. Cobb, Jr.	31
Figure 18. Sonic velocities for Ross #1, El Paso Natural	32
Figure 19. Sonic velocities for Stremler #1, Can-American Petrol., Ltd.	33
Figure 20. Densities for Montesano #1-X, El Paso Production	34
Figure 21. Densities for Kerry BN #34-11, L B Petroleum	35
Figure 22. Densities for Amoco-Weyerhaeuser #1-29, Amoco Production	36
Figure 23. Densities for Black Diamond #4-13, Voyager Petrole	37
Figure 24. Densities for Socal-Schroeder #1, Standard Oil of Calif.	38
Figure 25. Densities for Kingston #1, Mobil Oil	39
Figure 26. Densities for Soleduck #1, Eastern Petroleum Serv.	40
Figure 27. Densities for Socal Whidbey #1, Standard Oil of Calif.	41

Figure 28.	Densities for Squalicum Lake #1, Washuta Drilling & Company	42
Figure 29.	Calculated two-way travel time for Kerry BN #34-11, L B Petroleum	43
Figure 30.	Calculated two-way travel time for Amoco-Weyerhaeuser #1-29, Amoco	44
Figure 31.	Calculated two-way travel time for Blessing Siler #1, Carr Edward J	45
Figure 32.	Calculated two-way travel time for Washington State #1, Phillips	46
Figure 33.	Calculated two-way travel time for Brandt #1, McCroskey A E Syndicate	47
Figure 34.	Calculated two-way travel time for Brandt #2, McCroskey A E Syndicate	48
Figure 35.	Calculated two-way travel time for KSD #1, Geothermal Resources Int'l.	49
Figure 36.	Calculated two-way travel time for Socal-Schroeder #1, Standard Oil	50
Figure 37.	Calculated two-way travel time for Kingston #1, Mobil Oil	51
Figure 38.	Calculated two-way travel time for Pope & Talbot #18-1, Union Oil.	52
Figure 39.	Calculated two-way travel time for Socal Whidbey #1, Standard Oil	53
Figure 40.	Calculated two-way travel time for Pope & Talbot #3-1, Standard Oil	54
Figure 41.	Calculated two-way travel time for Dungeness #1, Standard Oil	55
Figure 42.	Calculated two-way travel time for R D Merrill Co. #1, Texaco Inc.	56
Figure 43.	Calculated two-way travel time for Silvana Community #12-1, Standard Oil	57
Figure 44.	Calculated two-way travel time for Merrill-Ring #1, Russell A. Cobb, Jr.	58
Figure 45.	Calculated two-way travel time for Ross #1, El Paso Natural	59
Figure 46.	Calculated two-way travel time for Stremler #1, Can-American Petrol., Ltd.	60

## INTRODUCTION

We describe sonic velocity and density log data from oil test wells being used to help develop a three-dimensional velocity model for the Puget Sound. These 3-D models will be used to calculate synthetic seismograms to help understand the lateral variations of strong ground motions in the Puget Lowland urban corridor. We present data from 29 sonic and density logs from 23 oil test wells to categorize the sonic velocities and densities of Cenozoic sedimentary basins in western Washington, primarily in Puget Lowland (Fig. 1).

The locations, elevations, and depths of the oil test wells, as well as the lease name, well number, operator, and completion year are presented in Table 1. In this table the wells are ordered by latitude from south to north. This information is taken from the Well History Control System (WHCS) One-line File, an on-line digital well-log database leased from Petroleum Information by the USGS Office of Energy Resources at Denver. Because the logs were run over a 28-year interval between 1958 and 1985, Table 1 provides information on the type of sonic and density tool used to make the log, as well as the other tools which were run simultaneously with these tools (normally caliper, spontaneous potential, and gamma-ray). Several of the sonic logs were made with older, short tools, with short spans between the source and receivers. Stratigraphic control is available for many of the wells [MacFarland, 1983; Brown and Ruth Laboratories, Inc., 1984; Johnson and others, 1996; Rau and Johnson, in press; S.Y. Johnson, written commun., 1998]. We note here that the formation picks for the Dungeness Spit #1 well (shown in Table 3 and on Figure 14) are inferred from the formation tops at the Dungeness Unit #1-54 well, located approximately 9 km to its southeast (T30N, R4W, Section 17; McFarland, 1983).

Where known (Table 2), the generalized stratigraphy encountered in the holes may be characterized as follows [MacFarland, 1983; Rau and Johnson, in press]. Tertiary sediment rocks (generally Eocene), underlie Pleistocene glacial deposits (clays, sands, and gravels), and in a few wells are in turn underlain by Eocene Crescent Formation volcanics (or other volcanics).

A map showing thickness of unconsolidated deposits (Jones, 1994) was used to help ascertain thicknesses of Pleistocene deposits at wells for which we lacked other stratigraphic information.

## WELL LOG ANALYSIS

Sonic and density logs were hand digitized at non-uniform intervals between 3 and 30 m to capture the significant variations of the logs with depth for frequencies up to 2 Hz. The sampling interval was adequate to estimate linear trends in the data over these intervals. We note that our sampling interval was not intended and is not sufficiently dense for the calculation of high-frequency (say >10 Hz) synthetic seismograms. For higher-frequency synthetics, it will be necessary to redigitize the logs with a finer sampling interval.

For the sonic logs, we picked transit times ( $\Delta t$ /ft) as a function of depth down the well. For the gamma-gamma density logs, we picked bulk density in  $\text{g/cm}^3$  as a function of depth down the well. For the neutron density porosity logs, we converted the logged density porosity ( $\Delta$ ) back to formation density ( $\rho_{fd}$ ) using  $\rho_{fd} = \rho_m + (\rho_f - \rho_m)\Delta$ , where the matrix density  $\rho_m = 2.65 \text{ g/cm}^3$ , and the fluid density  $\rho_f = 1.0 \text{ g/cm}^3$  [Ellis, 1987]. All of the logs analyzed here are plotted at a scale of 30.49 m = 5 cm (100 feet = 2 inches). Depths are measured from an arbitrary reference datum, normally the Kelley Bushing (KB), located 3.65 m (12 feet) above ground level. The downhole depths reported here have not been corrected for this small upward shift. Cased intervals of the wells and sections identified on the logs as having cycle skipping problems were not digitized. In some cases data from the logs were ignored: these data were associated with washouts, thick mudcake, invasion of drill fluids or large deviations from the general trend of density and sonic values having very limited depth extent, generally less than a few tens of feet [Ellis, 1987]. The digitized sonic log data were converted from transit times to velocities (m/s) and depths from feet to meters for both the sonic and density logs. Plots showing seismic velocities and densities as a function of depth for each well are presented in Figures 2 to 28. Although we digitized all repeated passes of tools in sections of the wells, we do not show these redundant passes in Figures 2 to 28.

We have calculated two-way travel times from the sonic logs and displayed these in Figures 29 to 46. This calculation required us to extrapolate sonic velocities to the surface. In some cases, the distance over which the first measured sonic velocity was judged to be too great to extrapolate it to the surface. In these cases, we calculated two-way travel times starting at the top of the sonic log.

### **COMMENTS ON THE WELL LOG DATA**

Although the deepest boreholes for which logs are included in this completion are more than 3.9 and 4 km deep respectively (Table 1), these deep wells were unfortunately not logged to the completed depth of the holes. The four longest sonic well logs are from the following wells (numbers in parens provide the logged intervals): Washington State #1 (300-3500 m), Socal Schroeder #1 (100-2950 m), Silvana Community #12-1 (150-2250 m), and the Socal Whidbey #1 (50-2050 m). The density logs from the same holes are generally slightly shorter. The four longest density well logs are from the following wells: Socal-Schroeder #1 (350-2950 m), Montesano 1-X (50-2100 m), Black Diamond #4-13 (200-2000 m), and Socal Whidbey #1 (300-2050 m).

The Amoco Weyerhaeuser #1-29 well was drilled and logged twice. The longest logs shown in this report are from the initial borehole; the second, shorter logs are from the first redrill. The sonic velocities measured in both wells provide comparable estimates in the overlap region (Fig. 3), however, there is a significant offset of  $0.2 \text{ g/cm}^3$  between the densities determined at the adjacent wells (Fig. 22). The origin of this mismatch is not currently known.

We have superimposed the formation tops from Table 2 on the well logs shown in this report. Often, but by no means always, these formation changes correspond to significant changes in the sonic velocity and density, and/or in the character of these logs. In Table 3 we show average velocities and linear regressions over intervals of the boreholes corresponding to known or inferred formations. The average sonic velocities represent the inverse of the average

transit times (slownesses) over the interval. In some cases, we show these averages and linear regressions for the entire log. As anticipated, average sonic velocities (generally 1450 to 2200 m/s) and densities ( $2.09 \text{ g/cm}^3$ ) in the Pleistocene glacial deposits are significantly lower than those of underlying Tertiary sedimentary rocks (Table 3). There is typically a general increase with depth in the sonic velocity within these Tertiary sedimentary rocks without a major change at the formation boundaries; although the low  $R^2$  values found for most of the linear regressions indicate that there is much scatter in this trend. The top of the Crescent Formation volcanics, however, does typically represent a large (over 1 km/s) increase in sonic velocity from the overlying Tertiary sedimentary rocks (Table 3).

Similar averages and linear regressions on the density log data are shown in Table 4. The density logs typically show even less evidence for contrasts between the Tertiary sedimentary rocks. There is, however, a significant ( $0.17 \text{ g/cm}^3$ ) contrast between these units and the underlying Crescent Formation volcanics at the Socal Whidbey #1 well.

The density log from Socal Schroeder #1 shows an apparent reversal within the Blakeley Formation, and densities of the andesitic volcanic rocks are lower than those of the Blakeley Formation (Table 4). This reversal is not accompanied by a similar reversal in sonic velocity, which, in combination with the generally poor quality of the density log for this well, suggest that the apparent reversal in density may be an artifact.

## **DATA AVAILABILITY**

The picks of density and seismic velocity shown in Figures 2 to 28 are available in Excel5 spreadsheets using anonymous ftp. The anonymous ftp address is: [andreas.wr.usgs.gov](ftp://andreas.wr.usgs.gov). Change the directory (cd) to /pub/outgoing/puget. The files are named `puget.sonic.xls.bin`, `puget.density.xls.bin`, and `puget.TT.xls.bin`, all in Mac Binary II format. Table 1 of this report in Excel5 format is also in this ftp site, labeled as `Table 1.bin`. Figure 1 is in Adobe Illustrator 6 format in file `OFR.Fig1`.



## DISCUSSION

Published compilations of S-wave data for sedimentary rocks provide some guidance for typical  $V_p/V_s$  ratios in Tertiary sedimentary units. Ohta et al. [1977] published  $V_s$  and  $V_p$  in two deep holes sampling Tertiary sediments in Japan. Hamilton [1979] compiled these and similar measurements in Russia to obtain  $V_p/V_s$  ratios in the uppermost 1 km of section within Tertiary sedimentary units. Finally, Castagna et al. [1985] compiled  $V_s$  and  $V_p$  data from S-wave and P-wave logging and other measurements to obtain  $V_p/V_s$  for a variety of clastic silicate rocks. Castagna et al. [1985] suggest that  $V_p/V_s$  ratios in sandstones reach an average value near 1.7 beginning at depths of about 2 km, whereas the  $V_p/V_s$  in noncalcareous shales reach an average value of about 2 at depths near 3 km.

## ACKNOWLEDGEMENTS

Zenon Valin, USGS, kindly performed a search of a digital database providing well locations and other well information. Larry Beyer, USGS, made many useful suggestions for digitizing the well log data. Sam Johnson, USGS, provided preprints of his papers on the biostratigraphy of many of the wells in Puget Sound, and reviewed an earlier version of this report. Weldon Rau and Bill Lingley, Washington Department of Natural Resources, provided stratigraphy for several of the boreholes.

This work was supported by the National Earthquake Hazards Reduction Program.

## REFERENCES CITED

- Brown and Ruth Laboratories, Inc., 1984, Regional petroleum geochemistry of Washington and Oregon: Brown and Ruth Laboratories, Inc., 1 v., looseleaf.
- Castagna, J.P., Batzle, M.L., and Eastwood, R.L., 1985, Relationships between compressional-wave and shear-wave velocities in clastic silicate rocks, *Geophysics*, v. 50, p. 571-581.
- Ellis, D.V., 1987, *Well Logging for Earth Scientists*, Elsevier, New York, 532 p.
- Gard, L.M., Jr., 1968, Bedrock geology of the Lake Tapps quadrangle, Pierce County, Washington: U.S. Geological Survey, Professional Paper 388-B, p. B1-B33.
- Hamilton, E.L., 1979,  $V_p/V_s$  and Poisson's ratios in marine sediments and rocks, *Journal of the Acoustical Society of America*, v. 66, p. 1093-1101.
- Johnson, S.Y., 1993, Analysis of Cenozoic subsidence at three sites in vicinity of the Seattle basin, Washington: U.S. Geological Survey Open-File Report 93-332, 17 p.
- Johnson, S.Y., Potter, C.J., Armentrout, J.M., Miller, J.J., Finn, C., and Weaver, C.S., 1996, The southern Whidbey Island fault, western Washington-An active structure in the Puget Lowland, Washington: *Geological Society of America Bulletin*, v. 108, p. 334-354, and oversized insert.

- Johnson, S.Y., Tennyson, M.E., Lingley, W.S., and Law, B.E., 1993, Petroleum geology of the State of Washington: U.S. Geological Survey, Professional Paper 1582, 40 pp.
- Jones, M.A., 1994, Thickness of unconsolidated deposits in the Puget Sound Lowland, Washington and British Columbia, U.S. Geological Survey, Water Resources Investigations Report (WRIR) 94-4133.
- McFarland, C.R., 1983, Oil and gas exploration in Washington, 1900-1982, Washington (State), Dept. of Natural Resources, Div. Geology and Earth Resources, Information Circular 75, 119 p.
- Ohta, Y., N. Goto, K. Shiono, H. Takahashi, F. Yamamizu, and S. Kurihara, 1977, Shear wave velocities in deep soil deposits; measurement in a borehole to the depth of 3500 meters and its significance, Jishin Gakkai (Seismological Society of Japan), v. 30, p. 415-433.
- Rau, W.W., and S.Y. Johnson, in press, Well stratigraphy and correlations, western Washington and northwest Oregon, U.S. Geological Survey, Map I-2621.

#### ABBREVIATIONS USED IN TABLE 1:

BHC - Borehole Compensated Sonic Log

CNFD - Compensated Neutron Formation Density\*

CFD gg - Compensated Formation Density (gamma-gamma)\*

CNF (ds) - Compensated Neutron Formation Density (Dual Spaced)\*

Cal. - Caliper

SP - Spontaneous Potential

GR - Gamma Ray

T3R3R - Sonic tool spacing (in feet) between transmitter (T) and receivers (R)

\*All density logging tools employ the backscattered gamma-ray technique, commonly called "gamma-gamma". The different names used here are either from different vendors or from different generations (having different trademark names).

Table 1. Oil Test Well Data.

Leasename	No. Operator	Field	Latitude	Longitude	Depth,		Elev,		Datum	T	R	S	Year	Sonic Log	Density Log	Other Logs
					ft	m	ft	m								
KERRYN BN	34-11 L B PETROLEUM	Carbonado	47.08853	-122.02711	5000	1524	1342	409	GL N 19 E 6	34	1985	BHC 5"	FDC/CNL	Cal., GR, SP		
AMOCO-WEYERHAEUSER	1-29 AMOCO PROD	Caldwell Creek	47.10136	-123.70526	13250	4040	390	119	GL N 19 W 8	29	1985	BHC 2"	FDC/CNL	Cal., GR		
BLESSING SILER	1 CARR EDWARD J	Bonney Lake	47.17198	-122.10001	7562	2305	658	201	GL N 20 E 6	31	1962	T3R3R		SP?		
WASHINGTON STATE	1 PHILLIPS PETROLEUM CO.	Lake Tapps	47.18277	-122.10367	12920	3939	625	191	GL N 20 E 5	36	1963	T3R3R		Cal., SP		
BRANDT	1 MCCROSKY A E SYNDICATE	Kummer Anticline	47.25790	-122.01697	3944	1202	657	200	KB N 21 E 6	34	1961	T3R3R				
BRANDT	2 MCCROSKY A E SYNDICATE	Kummer Anticline	47.25887	-122.01749	3411	1040	629	192	GL N 21 E 6	34	1961	T3R3R		SP		
BLACK DIAMOND	4-13 VOYAGER PETROLE	Kummer Anticline	47.32875	-121.91422	7270	2216			N 21 E 7	4	1983		Den			
KSD	1 GEOTHERMAL RESOURCES INTL	Kent	47.41684	-122.16891	9291	2833	438	134	GL N 22 E 5	4	1967	T3R2R		Cal., SP		
SOCAL-SCHROEDER	1 STANDARD OIL OF CALIF.	Brier-Puget Sound	47.79410	-122.26332	9675	2950	366	112	GL N 27 E 4	26	1972	BHC	CFD <sub>eg</sub> , CNFD	Cal., GR, SP		
KINGSTON	1 MOBIL OIL	Wildcat	47.80862	-122.49829	8648	2637	283	86	GL N 27 E 2	26	1972	T3R2R	CFD <sub>eg</sub>	Cal., GR, SP		
POPE & TALBOT	18-1 UNION OIL OF CALIF	Port Gamble	47.83406	-122.5892	4019	1225	190	58	GL N 27 E 2	18	1972	T3R3R		Cal., SP		
SOCAL WHIDBEY	1 STANDARD OIL OF CALIF.	Whidbey Island	47.97520	-122.40491	6693	2041	446	136	GL N 29 E 3	27	1972	T3R2R	CFD <sub>eg</sub>	Cal., GR, SP		
POPE & TALBOT	3-1 STANDARD OIL OF CALIF.	Whidbey Island	48.05327	-122.55345	4375	1334	318	97	GL N 30 E 2	28	1962	T3R3R		Cal., SP		
ENGSTROM	1 STANDARD OIL OF CALIF.	Dungeness Area	48.09395	-122.58786	7353	2242			N 30 E 2	17	1958	Son				
DUNGENESS	1 STANDARD OIL OF CALIF.	Pysht Area	48.16671	-123.13582	5105	1556	0	0	GL N 31 W 4	24	1965	T3R3R		Cal., GR		
R D MERRILL CO.	1 TEXACO INC	Silvana	48.19199	-124.09622	8460	2579	240	73	GL N 31 W 11	10	1966	T3R3R		Cal., SP		
SILVANA COMMUNITY	12-1 STANDARD OIL OF CALIF.	Pillar Point	48.19441	-122.23065	7419	2262	31	9	GL N 31 E 4	12	1958	T3R3R		SP		
MERRILL-RING	1 RUSSELL A. COBB, Jr.	Bellingham	48.21706	-124.13037	8519	2597	155	47	GL N 31 W 11	4	1960	TR1R		SP		
SQUALICUM	1 WASHUTA DRILLING &	Wildcat	48.77538	-122.35051	6180	1884	538	164	GL N 38 E 4	18	1968		CFD <sub>eg</sub>	Cal.		
ROSS	1 EL PASO NATURAL	Lynden	48.77798	-122.33031	4707	1435	695	212	GL N 38 E 4	17	1962	T3R3R		Cal., SP		
STREMLER	1 CAN-AMER PETROL., LTD		48.97983	-122.43848	8342	2543	122	37	GL N 40 E 3	4	1962	T3R3R		Cal., SP		

**Table 2. Stratigraphy in oil test wells**

Well Name	Formation	Depth (ft)	Depth (m)
Montesano #1-X*	Pleist. glacial		
	Top/Astoria Fm. (?)	250	76.2
	Top/Lincoln Creek Fm.	900	274.4
	Top/Humtulpis Fm.	6100	1859.8
Kerryn BN #34-11	Carbonado Fm.	0-4920	0-1500
Weyerhaeuser #1-29*	in Astoria (?) Fm.	1700	518.3
	Top/Lincoln Creek Fm.	3200	975.6
Blessing Siler Community #1	Puget Group?	820-7544	250-2300
Washington State #1□	Pleistocene	0-850	0-259
	Top/Spiketon Fm.	7000	259
	Top/Northcraft Fm.	7650	2134
	Top/Carbonado Fm.	12920	2332
Brandt #1	Puget Group	984-4018	300-1225
Brandt #2	Puget Group	656-3444	200-1050
Black Diamond #4-13	Quaternary Alluvium	0	0
	Lower Puget Group	50	15
	Raging River Fm.		
	regressive ss facies	1500	457
	deeper water facies	4500	1372
	transgressive ss facies	5500	1677
	Basement?	6500	1982
KSD #1 □□□	Renton Fm.	to 6870	to 2095
	Top/Tukwillia Fm.	6870	2095
	Top/Tiger Mountain Fm.	8700	2652
Schroeder #1*	in Pleist. glacial		
	Top/Blakeley Fm.	680	207.3
	Top/Renton Fm.	5660	1725.6
	Top/Mt. Perris volc. rx.	8000	2439.0
	Top/Puget Group	8850	2698.2
Kingston #1*	in Pleist. glacial	500	152.4
	Top/Blakeley Fm.	1720	524.4
	Top/Puget Group correlative?	3500	1067.1
	Top/Scow Bay ss	4800	1463.4
Pope & Talbot #18-1	Pleistocene	0	0

	Top/Blakeley Fm.	1328	405
	Top/Crescent Fm.	3369	1027
Soleduck #1 , □□□	Ozette Melange (Eocene and Miocene broken fm. and melange)	328-1066	100-325
Socal Whidbey #1*	in Pleist. glacial	100	30.5
	Top/Blakeley Fm.	600	182.9
	indurated ss	6000	1829.3
	Top/Crescent Fm. - Volcs.	6100	1859.8
	Top/Scow Bay ss	6500	1981.7
Pope & Talbot #3-1**	Pleistocene	0	0
	Top/Blakeley Fm. (upper part)	672	205
	Top/Blakeley Fm. (lower part)	2230	680
	Top/ss of Scow Bay/unnamed strata	3411	1040
Dungeness Unit #1-54 ,***	Pleist. glacial	0	0.0
	Top/Pleist. conglomerate	1560	476
	Top/Twin River Fm.	3282	1000.6
	Top/Crescent Fm. - Volcs.	6365	1940.5
Silvana Community #12-1*	in Pleist. glacial	520	158.5
	Top/Bulson Creek Fm.	900	274.4
	Top/Basaltic volc. rocks	7350	2240.9
Merrill-Ring #1□□□	Upper and Middle Twin River Group	to 5000	to 1524
	Middle Twin River Group	5000	1524
Squalicum Lake #1	Chuckanut Fm.	1886-5576	575-1700
Ross #1	Pleistocene	0	0
	Top/Chuckanut Fm.	843	257
Stremler #1	Chuckanut and Huntingdon Fm.	2542-7380	775-2250

\*Rau and Johnson (in press).

Field notes of Howard Gower.

Gard (1968).

Written communication from S.Y. Johnson, 1998.

□ Brown and Ruth Laboratories, Inc. (1984).

Proprietary industry report.

□□□ W. W. Rau, Washington State Dept. of Natural Resources, pers. comm., 1998.

\*\*Johnson and others (1996).

\*\*\*From MacFarland (1983).

**Table 3. Linear regression of sonic velocities in oil test well logs**

Wellname and number	Formation Name	Depth Interval (m)	Average Velocity (km/s)	Intercept Velocity (km/s)	Velocity Gradient (km/s/km)	R <sup>2</sup>
Kerryn BN #34-11	Carbonado Fm.	96-1481	4.36	3.83	0.84	0.32
Weyerhaeuser #1-29	All	532-1370	2.40	2.14	0.30	0.10
Weyerhaeuser #1-29 redrill	Lincoln Creek Fm.	1059-1414	2.62	3.90	-1.01	0.10
Blessing Siler #1	Puget Group?	229-2304	3.68	3.39	0.33	0.10
Washington State #1	Puget Group	303-3472	3.74	2.91	0.54	0.52
Brandt #1	Puget Group	421-1199	3.49	3.17	0.52	0.05
Brandt #2	Pleistocene	149-202	1.45			
	Puget Group?	229-1037	3.31	2.55	1.49	0.23
KSD #1	Puget Group	>2350	4.44	4.04	0.18	0.00
Socal-Schroeder #1	Pleist. deposits	0-207	1.59			
	Blakeley Fm.	207-1726	3.26	2.36	0.88	0.46
	Renton Fm.	1726-2439	3.30	5.70	-1.09	0.12
	Andesitic volc. rocks	2439-2698	3.78			
	Puget Group	>2698	3.54			
Kingston #1	Pleist. deposits	175-524	1.83	1.57	0.92	0.37
	Eocene sed. rocks	>524	2.88	2.04	0.85	0.44
Pope&Talbot #18-1	Pleistocene	0-405	N/A			
	Blakeley Fm.	405-1027	2.78	2.03	1.08	0.51
	Crescent Fm. volc.	1027-1225	4.65			
Socal-Whidbey #1	All	47-2040	2.69	1.80	1.09	0.78
	Pleist. deposits	31-183	1.71			
	Blakeley Fm.	183-1829	2.77	2.01	0.86	0.73
	Indurated Ss.	1829-1860	3.30			
	Crescent Fm. volc.	1860-1982	5.04			
	Scow Bay Ss.	>1982	4.25			
Pope&Talbot #3-1	All	165-1324	2.70	1.62	1.70	0.84
	Pleistocene	0-205	1.57			
	Blakeley Fm.	205-680	2.43	2.09	0.84	0.27
	Blakeley Fm.	>680	2.98	1.26	1.96	0.75
	Crescent Fm. volc.	>1040	3.83			
Dungeness #1	All Pleistocene?	236-850	2.28	1.97	0.70	0.09
	Pleist. deposits?	0-625	2.14	2.31	-0.28	0.01
	Pleist. conglomerates?	625-825	2.50			
	Twin River Fm.?	825-1500	2.59	1.73	0.74	0.50
	Crescent Fm.?	>1500	3.70			
R.D. Merrill #1	All	302-1829	2.50	1.94	0.61	0.54
Silvana Comm. #12-1	All	160-2255	3.04	2.38	0.62	0.47
	Pleist. deposits	159-274	2.17			
	Bulson Creek Fm.	274-2240	3.12	2.54	0.50	0.37
	Basaltic volc. rocks	>2240	4.87			
Merrill-Ring #1	All	1465-2271	2.50	-0.69	1.80	0.58
Ross #1	Pleistocene	0-257	1.92			
	Chuckanut Fm.	257-620	3.60	2.48	2.90	0.36
Stremler #1	Chuckanut and Huntingdon Fm.	765-2233	4.31	4.22	0.09	0.01

**Table 4. Linear regression of densities in oil test well logs**

Wellname and number	Formation Name	Depth Interval (m)	Average Density (g/cc)	Intercept Density (g/cc)	Density Gradient (g/cc)	R <sup>2</sup>
Montesano #1-X	All	59-2112	2.16	2.04	0.10	0.44
	Astoria (?) Fm.	76-274	2.11			
	Lincoln Creek Fm.	274-1860	2.15			
	Humptulips Fm.	>1860	2.36			
Kerryn BN #34-11	Carbonado Fm.	101-1475	2.61	2.57	0.05	0.08
Weyerhaeuser #1-29	All	532-1384	2.05	2.13	-0.08	0.06
Weyerhaeuser #1-29 redrill	Lincoln Creek Fm.	1062-1417	2.28	2.71	-0.30	0.46
Black Diamond #4-13	All	205-1985	2.53	2.59	-0.06	0.07
Socal-Schroeder #1	Pleist. deposits	0-207	N/A			
	Blakeley Fm.	207-1726	2.49			
	Renton Fm.	1726-2439	2.23			
	Andesitic volc. rocks	2439-2698	2.19			
	Puget Group	>2698	2.17			
Kingston #1	Pleist. deposits	175-524	2.09	1.95	0.40	0.34
	Eocene sed. rocks	524-1550	2.29	2.07	0.20	0.43
Soleduck #1	Tertiary broken fm?	96-316	2.20	2.11	0.50	0.10
Socal-Whidbey #1	Pleist. deposits	31-183	N/A			
	Blakeley Fm.	183-1829	2.23	2.03	0.20	0.56
	Indurated Ss.	1829-1860	2.47			
	Crescent Fm. volc.	1860-1982	2.64			
	Scow Bay Ss.	>1982	2.49			
Squalicum #1	Chuckanut Fm.	610-1687	2.52	2.45	0.07	0.06




Figure 1. Map showing location of wells having logs analyzed in this report.



Kerryn-BN #34-11, L. B. Petroleum




Figure 2.

Weyerhaeuser #1-29, Amoco Production Company




Figure 3.

Blessing Siler Community #1, Edward J. Carr




Figure 4.

Washington State #1, Phillips Petroleum Company




Figure 5.

Brandt #1, A. E. McCroskey




Figure 6.

Brandt #2, A. E. McCroskey




Figure 7.

KSD #1, Geothermal Resource International, Inc.




Figure 8.

Socal-Schroeder #1, Standard Oil Co. of California




Figure 9.



Kingston #1, Mobil Oil Corporation




Figure 10.

Pope-Talbot #18-1, Union Oil Co. of California




Figure 11.

Socal Whidbey #1, Standard Oil Co. of California




Figure 12.

Pope & Talbot #3-1, Standard Oil Co. of California




Figure 13.

Dungeness Spit #1, Standard-Union




Figure 14.

R. D. Merrill Company #1, Texaco, Inc.




Figure 15.

Silvana Community #12-1, Standard Oil Co. of California




Figure 16.

Merrill-Ring #1, Russell A. Cobb, Jr.




Figure 17.



Ross #1, El Paso Natural Gas




Figure 18.

Stremler #1, Can American Petroleum, Ltd.




Figure 19.

Montesano I-X, El Paso Products Company




Figure 20.

Kerryn-BN #34-11, L. B. Petroleum




Figure 21.

Weyerhaeuser #1-29, Amoco Production Company




Figure 22.

Black Diamond #4-13, Voyager Petroleum




Figure 23.

Socal-Schroeder #1, Standard Oil Co. of California




Figure 24.

Kingston #1, Mobil Oil Corporation




Figure 25.



### Soleduck #1, Eastern Petroleum




Figure 26.

Socal Whidbey #1, Standard Oil Co. of California




Figure 27.

Squalicum Lake #1, Washuta Drilling Company




Figure 28.

Kerryn-BN #34-11, L. B. Petroleum




Figure 29.

Weyerhaeuser #1-29, Amoco Production Company




Figure 30.

Blessing Siler Community #1, Edward J. Carr




Figure 31.

Washington State #1, Phillips Petroleum Company




Figure 32.

Brandt #1, A. E. McCroskey




Figure 33,



Brandt #2, A. E. McCroskey




Figure 34.

KSD #1, Geothermal Resource International, Inc.




Figure 35.

Socal-Schroeder #1, Standard Oil Co. of California




Figure 36.

Kingston #1, Mobil Oil Corporation




Figure 37.

Pope-Talbot #1 8-1, Union Oil Co. of California




Figure 38.

Socal Whidbey #1, Standard Oil Co. of California




Figure 39.

Pope & Talbot #3-1, Standard Oil Co. of California




Figure 40.

Dungeness Spit #1, Standard-Union




Figure 41.



R. D. Merrill Company #1, Texaco, Inc.




Figure 42.

Silvana Community #12-1, Standard Oil Co. of California




Figure 43.

Merrill-Ring #1, Russell A. Cobb, Jr.




Figure 44.

Ross #1, El Paso Natural Gas




Figure 45.

Stremler #1, Can American Petroleum, Ltd.




Figure 46.